

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 39/395, 39/40, 39/42, 38/00, G01N 33/53, 33/537, 33/543, 33/563, C12N 7/00, 7/01, C07K 16/00, C12P 21/08	A1	(11) International Publication Number: WO 97/06823 (43) International Publication Date: 27 February 1997 (27.02.97)
(21) International Application Number: PCT/US96/13215 (22) International Filing Date: 15 August 1996 (15.08.96) (30) Priority Data: 60/002,462 16 August 1995 (16.08.95) US (71) Applicant: UNIVERSITY OF MASSACHUSETTS MEDICAL CENTER [US/US]; 55 Lake Avenue North, Worcester, MA 01655 (US). (72) Inventors: HERRMANN, John, E.; 10 Newton Street, Northboro, MA 01532 (US). BLACKLOW, Neil, R.; 144 Summer Street, Weston, MA 02193 (US). (74) Agent: FASSE, J., Peter; Fish and Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).		(81) Designated States: AU, CA, JP, KR, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: MONOCLONAL ANTIBODIES FOR DETECTING NORWALK VIRUS (57) Abstract Monoclonal antibodies and their use in assays to determine the presence of Norwalk virus, Norwalk virus particles, or Norwalk virus-specific IgM, in a biological sample. One assay includes the steps of contacting the sample with a monoclonal antibody that specifically binds to Norwalk virus particles in human stool; reacting the sample and the antibody for a time and under conditions that allow the formation of an immunocomplex between the antibody and any Norwalk virus in the sample; and detecting the immunocomplex, the presence of the immunocomplex indicating the presence of human Norwalk virus in the sample. Another assay is an IgM capture antibody immunoassay.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

MONOCLONAL ANTIBODIES FOR DETECTING NORWALK VIRUSBackground of the Invention

This invention relates to methods of detecting
5 Norwalk virus and Norwalk virus-specific IgM with
monoclonal antibodies.

Norwalk virus, the prototype calicivirus, is a
major cause of outbreaks of gastroenteritis, affecting
primarily older children and adults. Although Norwalk
10 virus has been known since its first outbreak in Norwalk,
Ohio in 1968, viral particles were not visualized until
1972. The 27 nm diameter particles were first observed
in a stool sample by immune electron microscopy using
serum from a symptomatic patient. Norwalk virus has not
15 been serially propagated in cell cultures, which has
inhibited the study and diagnosis of this virus and its
antigens. Instead, diagnosis is accomplished by PCR,
ELISA, antigen detection, electron microscopy, or by
serologic methods.

20 Diagnosis of Norwalk virus infections by
serological means or by antigen detection in the past has
relied on use of clinical materials, e.g., human serum,
obtained from volunteers infected with the virus. The
Norwalk virus genome has been cloned and a Norwalk virus
25 capsid protein has been expressed as described in Jiang
et al., Science, 250:1580-1583, 1990 and Jiang et al., J.
Virol., 66:6527-6532, 1992. Antisera to the recombinant
Norwalk virus (rNV) capsid protein have been prepared and
used in an ELISA for antigen detection. The ELISA was
30 highly specific for the prototype Norwalk virus, and
showed that some virus isolates that had been previously
identified as Norwalk-related viruses by tests using
human reagents were distinct from prototype Norwalk virus
(Graham et al., J. Infect. Dis., 170:34-43, 1994).
35 However, antisera to the rNV capsid protein have not been

- 2 -

demonstrated to react to any stool samples from naturally occurring Norwalk virus outbreaks.

Moreover, the use of IgG-based serologic methods for Norwalk virus diagnosis is limited because these
5 methods require the use of paired acute and convalescent phase sera. If only late-stage acute serum is available, showing a clear positive increase in Norwalk virus-specific IgG may not be possible if the acute phase serum already has a high titer.

10 Summary of the Invention

The invention features new monoclonal antibodies that specifically bind to Norwalk virus particles in stool from a human patient infected with Norwalk virus, and in particular, two new monoclonal antibodies
15 designated herein as 1C9 and 1D8.

In another aspect, the invention generally features immunoassays to determine the presence of Norwalk virus or virus antigens, e.g., particles, in a biological sample, e.g., a stool sample, by contacting
20 the sample with a monoclonal antibody of the invention, e.g., 1C9 or 1D8; reacting the sample and the antibody for a time and under conditions that allow the formation of an immunocomplex between the antibody and any Norwalk virus or virus antigens in the sample; and detecting the
25 immunocomplex, the presence of the immunocomplex indicating the presence of human Norwalk virus or virus particles in the sample. The viral antigen can be, for example, the 58 KDa Norwalk virus protein described herein.

30 In one embodiment of this assay, the immunocomplex can be detected by a competitive immunoassay by reacting the monoclonal antibody with the sample and with a competing antigen to which the monoclonal antibody is known to specifically bind, e.g., a labeled Norwalk virus

- 3 -

antigen or an immobilized competing antigen such as a recombinant Norwalk virus capsid protein. The competing antigen can be labeled or immobilized.

In another embodiment, the immunoassay is a sandwich immunoassay that uses a second antibody, e.g., a monoclonal antibody, that either also binds Norwalk viral particles or binds to the first monoclonal antibody, one of the two antibodies being immobilized and the other being labeled using standard techniques. In the sandwich immunoassay procedures, the Norwalk viral particle-binding antibody can be a capture antibody attached to an insoluble material, and the second Norwalk viral particle-binding antibody can be a detector or labeling antibody.

The immunoassay in another embodiment is a Norwalk virus antigen-based IgM-capture enzyme immunoassay (EIA) that uses a monoclonal antibody of the invention. This immunoassay enables the relatively early detection of Norwalk virus, and requires only one serum sample, compared to two samples (taken at about four week intervals) required for typical IgG seroconversion assays. This assay can be used to determine the presence of Norwalk virus-specific IgM in a biological sample, and includes the steps of isolating, e.g., immobilizing, IgM in the sample; contacting the isolated IgM with purified Norwalk virus or Norwalk virus particles for a time and under conditions that allow the formation of a Norwalk virus-IgM complex between any Norwalk virus-specific IgM in the sample and the purified Norwalk virus or Norwalk virus particles; contacting the Norwalk virus-IgM complex with a monoclonal antibody that specifically binds to Norwalk virus in human stool; reacting the complex and the monoclonal antibody for a time and under conditions that allow the formation of an immunocomplex between the monoclonal antibody and any Norwalk virus-IgM complexes;

- 4 -

and detecting the immunocomplex, the presence of the immunocomplex indicating the presence of Norwalk virus-specific IgM in the sample.

In this assay, the IgM can be isolated with an
5 anti-human IgM antibody, such as a rabbit anti-human IgM antibody.

The immunoassays can be used to determine Norwalk virus or virus particles in samples including stool, intestinal or duodenal aspirates, urine, or peritoneal,
10 serum, blood, and lymphatic or other bodily fluids. The assays are useful for diagnosis of individual cases of Norwalk virus infection, as well as epidemiological studies of Norwalk virus outbreaks.

In another aspect, the invention features anti-
15 idiotypic antibodies that specifically bind to the monoclonal antibodies of the invention, e.g., 1C9 or 1D8. These anti-idiotypic antibodies can be used in a vaccine suitable for administration to humans.

The invention also features immunoassay kits for
20 detecting Norwalk virus in human biological samples, comprising one or more Norwalk virus-binding monoclonal antibodies and the means for determining binding of the antibody or antibodies to Norwalk virus or virus particles in a biological sample. In preferred
25 embodiments, the kit includes one or both monoclonal antibodies 1C9 and 1D8, or the competing antigens described above.

By "purified" Norwalk virus or virus particles is meant virus or virus particles that are at least 60%, by
30 weight, free from the proteins and naturally-occurring organic molecules with which they are naturally associated. Preferably, the preparation is at least 75%, more preferably at least 90%, and most preferably at least 99%, by weight, Norwalk virus or virus particles.
35 Purified Norwalk virus particles include recombinantly

- 5 -

produced polypeptides or peptides, e.g., Norwalk virus capsid peptides. A purified Norwalk virus or virus particles can be obtained, for example, by affinity chromatography using antibodies or monoclonal antibodies
5 described herein, by physical purification techniques described herein, or by other standard techniques.

The term "Norwalk virus antigen," as used herein, includes antigenic proteins or polypeptides of the virus, either naturally occurring or recombinantly or
10 synthetically produced, to which antibodies of the invention specifically bind. The term thus includes the 58 KDa Norwalk virus protein described herein. The term "virus particle" is used to describe antigenic particles to which antibodies of the invention specifically bind.

15 By "specifically binds" is meant an antibody which recognizes and binds Norwalk virus or Norwalk virus particles or antigens, but which does not substantially recognize and bind other molecules in a sample, e.g., a biological sample, which naturally includes Norwalk virus
20 or virus particles.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and
25 materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are
30 incorporated by reference. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

Other features and advantages of the invention will be apparent from the following detailed description,
35 and from the claims.

- 6 -

Brief Description of the Drawings

Fig. 1 is a graph showing the titration of Norwalk virus antigen in a stool specimen from a volunteer by monoclonal antibody ELISA (closed circles) and polyclonal antibody (open circles). Points above the dotted line represent positive values for the immunoassays.

Fig. 2A is a photograph of a gel showing untreated, radiolabeled translation products of Norwalk virus ORF-2. Numbers on the left are molecular masses in kilodaltons.

Fig. 2B is a photograph of a gel showing immunoprecipitation of radiolabeled Norwalk virus ORF-2 translation products with Norwalk virus-specific monoclonal antibodies 1D8 (lane 1, 3 μ L reaction product immunoprecipitated; lane 2, 15 μ L) and 1C9 (lane 3, 3 μ L; lane 4, 15 μ L) compared to immunoprecipitation with astrovirus-specific monoclonal antibody 8E7 (lane 5, 3 μ L; lane 6, 15 μ L). Norwalk monoclonal antibodies 1D8 (lane 7) and 1C9 (lane 8) were also tested for reactivity with radiolabeled astrovirus ORF-2 translation products (capsid). Numbers on the right are molecular masses in kilodaltons.

Detailed Description

Applicants have obtained monoclonal antibodies that specifically bind Norwalk viral particles derived from a human stool sample from a patient infected with the virus. These antibodies are useful in various immunoassays to detect the presence of Norwalk virus or virus particles or Norwalk virus-specific IgM in clinical samples from outbreaks of Norwalk virus infection.

Purification of Norwalk Viral Particles

Norwalk virus was obtained from the diarrheal stool of a volunteer who had been inoculated with Norwalk

- 7 -

virus strain 8FIIa. Virus was partially purified from stool material by differential centrifugation and banding in CsCl gradients, essentially according to the procedures described by Greenberg et al., J. Virol., 5 37:994-999, 1981. Fractions were collected, assayed for density, and tested by ELISA for Norwalk virus antigen. Fractions which had densities of 1.34 to 1.40 and showed high reactivity by ELISA as described below were pooled, dialyzed against 0.01 M phosphate-buffered saline, pH 10 7.2, and used as virus inoculum for hybridoma production.

Hybridoma Production

Hybridomas were produced as described in Herrmann et al., J. Clin. Microbiol., 33:2511-2513, 1995. BALB/c mice (eight-week-old females) were inoculated 15 subcutaneously with 0.1 ml of stool-derived Norwalk virus emulsified in an equal volume of TiterMax adjuvant (Vaxcel, Inc., Norcross, GA). Mice were given a second subcutaneous inoculation three weeks later. The mice were given two more inoculations (intraperitoneal) of 20 Norwalk virus in Freund's incomplete adjuvant, and one intraperitoneal inoculation of virus without adjuvant, all two weeks apart. Five days after the last inoculation, mouse spleens were fused to SP 2/0 myeloma cells with Kodak polyethylene glycol 1450 plus DMSO, 25 according to the procedures described by Lane, J. Immunol. Methods, 181:223-228, 1985. Hybrid cells were seeded onto 24 well plates in HAT medium containing 10% Hybridoma Cloning Factor (IGEN, Inc., Rockville, MD). After seven days, hybridomas were screened for antibodies 30 which reacted with Norwalk virus antigen from stools obtained from volunteers.

Hybridomas that secreted such antibodies were cloned twice by use of the limiting dilution technique. Ascitic fluids for all clones were prepared in BALB/c

- 8 -

mice. The techniques described herein can be used to obtain monoclonal antibodies that specifically bind to Norwalk virus particles from stool samples such as the particular antibodies 1C9 and 1D8 described herein.

5 Immunoassays

Enzyme Immunoassays of Stool Samples

Enzyme immunoassays (EIAs) were used to screen hybridoma supernatant fluids and to test virus-specific reactivity as described in Herrmann et al., J. Clin.

- 10 Microbiol., 33:2511-13, 1995. For these tests, stool extracts known to contain Norwalk virus antigen, and rNV capsid protein were tested by modification of an EIA for detecting Norwalk virus which used polyclonal human sera described in Herrmann et al., J. Med. Virol., 17:127-133,
15 1985.

- For the indirect EIA used to screen hybridomas, wells of polyvinyl chloride microtiter plates were coated with pre- and post-challenge sera (four-week convalescent) from a volunteer infected with Norwalk
20 virus. The wells were coated for 24 hours at room temperature (20-22°C), and post-coated with 1% w/v bovine serum albumin in 0.01 M phosphate-buffered saline (PBS) for 24 hours at 4°C. Suspensions of stool-derived Norwalk virus or rNV capsid protein were added to the
25 wells and incubated 18-24 hours at 20-22°C. The plates were washed with PBS and 0.05 ml of hybridoma supernatant fluids diluted in 50% fetal calf serum and 50% 0.025 M Tris-HCl buffer (pH 7.2) with 0.015% Tween 20, were added and incubated for 1 hour at 37°C. Peroxidase-labelled
30 goat antibody specific for mouse IgG (Kirkegaard and Perry Laboratories, Gaithersburg, MD), at 1 µg/mL in the Tris buffer used above, was added and incubated for 1 hour at 37°C. The plates were washed five times with PBS, soaked for 30 seconds with PBS containing 0.05%

- 9 -

Tween 20, and washed again. Substrate for peroxidase (0.05 mL, O-phenylene-diamine- H_2O_2 ; Abbott Laboratories, North Chicago, IL) was added for 10 minutes, and the reaction was stopped with 0.1 mL of 1 N H_2SO_4 .

5 Biotin/avidin labeling systems can also be used.

The absorbance of the solution was measured at 492 nm in a plate-reader spectrophotometer. After monoclonal antibodies were obtained, a direct EIA was used for testing stool samples; however, other standard
10 immunoassays can also be used. The same monoclonal antibodies were used for coating plates and for antigen detection. The antibodies were purified from ascitic fluid by ammonium sulfate precipitation, and either used directly for coating plates or labelled with peroxidase
15 for use as detector antibodies. The antibodies were labelled with peroxidase by the periodate method of Wilson and Nakane, "Recent developments in the periodate method of conjugating horseradish peroxidase (HRPO) to antibodies," p. 215-224. In: Knapp et al. (eds.),
20 Immunofluorescence and Related Staining Techniques (Elsevier/North Holland Biomedical Press, Amsterdam, 1978).

For the EIA, wells were coated with monoclonal antibodies (2 $\mu\text{g/mL}$) for 18 to 24 hour at 20-22°C,
25 washed, and post coated with Superblock (Pierce Chemical Co., Rockford, IL) 4 to 6 hours. The plates were washed, and stool samples or controls (known positive and negative stool samples) added and incubated for 18 hours at 20-22°C. Plates were washed, and the peroxidase
30 conjugated antibodies added and incubated at 37°C for 2 hours. The remaining procedures were as described for the indirect EIA. Samples were considered positive for monoclonal antibody to Norwalk virus in the screening tests, or for Norwalk virus antigen in the virus-
35 specificity tests, if the absorbance value was both ≥ 0.1

- 10 -

and three or more times the negative control (wells coated with pre-immune serum in the screening tests, and wells coated with an unrelated monoclonal antibody in the virus specificity tests).

5 Antigen-based IgG-capture EIA of Serum Samples

Recombinant Norwalk virus antigen-based IgM-capture enzyme immunoassays were used to test serum samples from two groups of human volunteers infected with Norwalk virus, and who had either an IgG antibody
10 increase against recombinant NV capsid protein, or were virus positive.

Volunteer Group I samples consisted of pre-inoculation sera followed by sera collected from five days to 22 weeks post-inoculation of the 8FIIa strain of
15 Norwalk virus (these samples were collected in the 1970s and stored at or below -20°C). Volunteer Group II was inoculated with 8FIIa in 1995, and samples consisted of pre-inoculation sera followed by sera collected at 4, 5, 8, 14 and 21 days post-challenge. Samples from two
20 natural Norwalk virus outbreaks were also tested. The first set of outbreak samples (Outbreak Group I) was from a Norwalk virus clam-associated outbreak of gastroenteritis in Hawaii in 1983. The second set of outbreak samples (Outbreak Group II) was from an outbreak
25 of gastroenteritis in Erie County, New York, in 1986.

Sera from volunteers challenged with Hawaii virus and sera from a Snow Mountain virus outbreak were also tested, in addition to sera from a group of normal human adult donors from a hospital blood bank and non-
30 symptomatic children admitted to the hospital for reasons other than gastroenteritis. All sera were stored at or below -20°C.

For the IgM capture antibody ELISA, polyvinyl plates (Dynatech Laboratories, Inc., Chantilly, VA) were
35 coated with unlabeled rabbit anti-human IgM (Fc5μ)

- 11 -

(Accurate Chemical, Westbury, NY) at a concentration of 0.25 µg/50 µl of 0.1 M PBS per well. Plates were incubated at 37°C for 2 hours, washed three times with PBS plus 0.15% TWEEN-20^m and blocked overnight with 5.0% BSA and 0.25% Bloom 60 gelatin in PBS. The plates were washed three times and duplicate two fold serial dilutions of human serum were made in 50% fetal bovine serum (50% 0.025 M Tris-HCL buffer (pH 7.2) with 0.015% TWEEN-20^m), and incubated for 1 hour at 37°C.

The plates were washed five times and 50 ng of recombinant Norwalk virus (rNV) capsid antigen, in 50 µl of the Tris-HCL-FBS buffer described above, was added to each well of one of the duplicate rows. To the second row, 50 µl of Tris-HCL-FBS buffer without rNV was added.

After overnight incubation at 20-22°C, the plates were washed five times and a combination of the two murine monoclonal antibodies 1C9 and 1D8 was added in Tris-HCL-FBS buffer to all wells and incubated at 37°C for 1 hour. The plates were washed and peroxidase-labeled goat anti-mouse immunoglobulin G (Kirkegaard & Perry Laboratories, Inc., Gaithersburg, MD) at 1 µg/ml in Tris HCL-FBS buffer plus 1% normal rabbit serum was added and incubated for 1 hour at 37°C. The plates were washed five times. Substrate for peroxidase (0.05 ml of 0-phenylenediamine-H₂O₂, Abbott Laboratories, North Chicago, IL.) was added and left for up to 10 minutes. The reaction was stopped with 0.1 ml of 1 N H₂SO₄.

The absorbance (optical density) of the solution was measured at 492 nm (A₄₉₂) in a plate-reader spectrophotometer. The endpoint titer was defined as the highest dilution of serum giving an optical density at A₄₉₂ of greater than or equal to 0.200 units higher than the corresponding antigen deficient well. To confirm IgM specificity, IgG and IgA were removed from the Group I

- 12 -

volunteer sera with Quik-SepIgM (Isolab Inc., Akron, OH) and tested in the same manner as the whole serum.

Radioimmunoprecipitation Tests

Radioimmunoprecipitation tests were done to
5 determine the reactivity of the monoclonal antibodies
with Norwalk virus capsid proteins produced in vitro.
These proteins were produced from full-length open
reading frame (ORF)-2 clones, which were described in
Jiang et al., J. Virol., 66:6527-6532, 1992, to code for
10 the Norwalk virus capsid protein. Reverse transcriptase-
PCR was used to obtain full-length ORF-2 clones from
Norwalk virus RNA isolated by standard silica extraction
from a fecal sample containing intact virus particles.
Oligo (dT) was used for reverse transcription to produce
15 first-strand cDNA.

Specific Norwalk virus primers were used in first
found PCR reactions. PCR products were amplified using a
"half-nested" approach (same 5'-end primers as in the
first round, with a unique 3'-end primer, internal to the
20 first round 3'-end primer). Second round PCR products
were made blunt-ended with T4 DNA polymerase and digested
with *Cla I*. The product (with a staggered 5'-end and a
blunt 3'-end) was ligated into *Cla I* and *Sma I*-digested
pBluescript KS-. ORF-2 was expressed in vitro in a
25 coupled transcription-translation system (Promega Corp.,
Madison, WI) and products were metabolically labeled with
[³⁵S]-methionine. Immunoprecipitations were performed as
previously described for astrovirus capsid proteins in
Lewis et al., J. Virol., 68:77-83, 1994.

30 The results of the immunoprecipitation studies are
given in Figs 2A and 2B. A predominant 58 kDa band,
indicative of Norwalk virus protein, was seen in crude
lysates. Norwalk virus-specific proteins were
immunoprecipitated with monoclonal antibodies 1D8 (lane

- 13 -

1, 2) and 1C9 (lane 3, 4). The predominant product seen in lanes 1 to 4 is a 58 kDa protein. An astrovirus-specific monoclonal antibody (8E7) failed to immunoprecipitate any Norwalk virus ORF-2 translation products (lane 5, 6). Conversely, Norwalk virus-specific monoclonal antibodies 1D8 and 1C9 did not immunoprecipitate any astrovirus capsid proteins (lanes 7, 8).

Testing of Monoclonal Antibodies

10 The selected hybridomas that produced antibodies reactive with both stool-derived Norwalk virus and rNV capsid protein were designated 1C9 and 1D8. No hybridomas were found that produced antibodies which reacted only with stool-derived virus or only with rNV capsid protein. Both antibodies were isotype IgG1. The two antibodies 1C9 and 1D8 appeared to be directed against different epitopes on the basis of blocking ELISA tests and the finding that a mixture of the two gave higher ELISA values than either one tested singly. This was confirmed by finding that only monoclonal antibody 1D8 reacted in an ELISA with a synthetic peptide to the N-terminus of the capsid. Based on these findings, a mixture of the two antibodies was used both for coating plates and as detector antibodies.

25 Stool samples containing other viruses were used for specificity testing of the monoclonal antibodies obtained. These include stools containing various U.K. strains of human calicivirus, and samples containing astroviruses, enteric adenoviruses, rotaviruses, Hawaii virus, Snow Mountain virus, unclassified "small round" viruses, and additional samples from Norwalk virus infected volunteers.

To determine the sensitivity of the monoclonal antibody ELISA compared to our previously developed

- 14 -

polyclonal ELISA, a volunteer stool was diluted in PBS and tested by both immunoassays. The results are shown in Fig. 1. The monoclonal antibody ELISA showed greater reactivity (higher A_{492} values) and 2-fold higher
5 sensitivity in detecting Norwalk virus in a diluted stool specimen than the polyclonal antibody ELISA. Both of the ELISAs shown are of the direct type. The detection limit for the monoclonal antibody ELISA, as determined by end-point titration of rNV capsid antigen, was 0.05
10 ng/microwell or 1 ng/ml, which is suitable for methods of detecting Norwalk virus particles in clinical samples.

Detection of Norwalk Virus Infection in Human Samples

Various immunoassay formats are suitable for the Norwalk virus particle detection methods of the
15 invention. For example, the ELISA, IgM-capture EIA, and radioimmuno-precipitation test described above can be used.

Human Stool Samples

In particular, representative types of
20 gastroenteritis viruses in stool samples from human volunteers previously found to contain Norwalk virus by a polyclonal ELISA as described in Herrmann et al., J. Med. Virol., 17:127-133, 1985, were used to test the specificity of the monoclonal antibodies for Norwalk
25 virus in an ELISA. The results, presented in Table 1, indicate that the monoclonal antibodies did not react with any normal stool samples or with samples containing viruses other than Norwalk virus. However, the monoclonal antibodies of the invention reacted
30 specifically to all 15 Norwalk virus samples tested that were previously found to be positive in a polyclonal antibody ELISA.

- 15 -

Table 1

	Patient group	No. of patients	No. ELISA positive
5	<u>Volunteers inoculated with</u>		
	Norwalk virus	15	15
	Hawaii virus	9	0
	<u>Patients infected with</u>		
10	Rotavirus	20	0
	Astrovirus	7	0
	Enteric adenovirus	7	0
	U. K. Calicivirus	5	0
	Snow Mountain virus	1	0
15	"Small round virus"	9	0
	Diarrhea of unknown etiology	30	0
	Stools from patients without diarrhea	25	0
20			

The reactivity of monoclonal antibodies with Norwalk virus in stools was determined primarily with stools obtained from volunteers infected with a strain of Norwalk virus circulating in 1968 (the "prototype" virus). Sera prepared against the recombinant Norwalk virus capsid protein have been shown to be highly specific for this prototype Norwalk virus (Graham et al., J. Infect. Dis., 170:34-43, 1994), and have not been shown to react with any stool samples from natural Norwalk virus outbreak stool samples.

It is believed that the monoclonal antibodies of the invention prepared against native virus from actual stool samples should be more broadly reactive than rNV capsid protein-generated antibodies with various Norwalk virus strains in naturally occurring outbreaks.

Strains from additional Norwalk virus outbreaks, especially recent ones, can be obtained and tested to

- 16 -

determine the breadth of reactivity of the monoclonal antibodies of the invention for diagnosis of Norwalk virus gastroenteritis caused by different strains of the virus.

5 Human Serum Samples

 All pre-challenge serum samples tested showed no detectable IgM. Norwalk virus-specific IgM was detected in nine of ten Volunteer Group I samples after virus inoculation (titers of 100 to 6400, 1/serum dilution).
10 IgM was detected as early as nine days after inoculation, but two samples of ten were negative five days after inoculation. IgM titers less than 25 were considered negative in all tests. Samples were taken at two different times from each volunteer, ranging from 5 days
15 to 7 weeks after inoculation for the first sample, and from 3 to 22 weeks after inoculation for the second sample.

 In Volunteer Group II, all five volunteers were IgM negative before and up to five days after exposure,
20 but all became IgM positive (1/serum dilution of 25 or higher) by eight days and were still positive 21 days after exposure. Samples were taken from each of the five volunteers at six time points: preinoculation, and 4, 5, 8, 14, and 21 days after inoculation.

25 In an rNV capsid antigen IgG assay, all of the subjects in volunteer Groups I and II showed Norwalk virus-specific rises in IgG titer. Sera from subjects in volunteer Group I that had IgG and IgA removed by Quik-Sep showed no significant differences in IgM antibody
30 titers as compared with untreated sera.

 Acute phase sera from the nine patients in Outbreak Group I (clam-associated outbreak in Hawaii in 1983) were collected from 6 to 8 days after onset of illness. Convalescent phase sera were collected 3 to 6
35 weeks post illness. All nine patients in Outbreak Group

- 17 -

I had high IgM antibody titers in both acute (800 to 12,800) and convalescent sera (3,200 to 25,600). Four patients showed four-fold rises in IgM levels. All nine patients had high IgG titers to rNV capsid antigen and
5 three had four-fold rises in IgG titer by EIA methods.

Outbreak Group II consisted of ten patients (gastroenteritis outbreak in Erie, NY in 1986), all of whom had four-fold or greater rises in IgG antibodies to rNV capsid protein. Seven of the subjects had acute sera
10 collected four days after a point source exposure and convalescent sera collected after two weeks post exposure. The other three subjects had acute sera collected between five and eight days post-exposure and convalescent sera collected three weeks post-exposure.
15 All of the Outbreak Group II subjects had acute sera lacking IgM antibodies except one whose exposure had been five to eight days earlier. Six of the nine persons whose acute phase sera were IgM negative seroconverted to Norwalk virus-specific IgM by the two or three week
20 samples. The one patient whose acute phase serum was IgM positive maintained a high IgM titer.

Three pairs of sera from subjects challenged with Hawaii virus and four pairs of sera from individuals in a Snow Mountain virus outbreak were tested for Norwalk
25 virus-specific IgM. All seven serum pairs were Norwalk virus IgM negative in both acute and convalescent phase samples although they cross-reacted in the rNV capsid antigen IgG assay. Eighty normal sera from subjects ranging from age one to age 59 were also tested. Four of
30 20 children aged 1 to 4, 12 of 20 children aged 5 to 17, 15 of 20 adults aged 18 to 29, and all 20 adults aged 30 to 59 were positive for IgG-specific Norwalk antibodies at a 1:800 serum dilution. On the other hand, none of the normal sera were IgM positive at a 1:25 serum
35 dilution for any age group.

- 18 -

Based on these studies, it appears that specific IgM is present only after recent exposure to Norwalk virus. The time that IgM specific antibodies can first be detected is important in defining NV outbreaks. In these studies, the earliest detection was at 5 days post inoculation, and all samples were positive by 8 or 9 days post inoculation. In Volunteer Group II, all five volunteers were negative at five days post inoculation and all were IgM-antibody positive by day eight. In outbreak Group I, all patients were highly positive for IgM by five to eight days post onset of illness. It appears from these groups that Norwalk virus-specific IgM develops by approximately eight days after virus exposure, and thus the capture IgM EIA described above is suitable for use in early detection of Norwalk virus infections.

Other Embodiments

Those skilled in the art will realize that known techniques can be used to raise anti-idiotypic antibodies, e.g., idiotypic antibodies that will bind to the monoclonal antibodies of the invention. Anti-Norwalk virus particle-binding antibodies can be used as immunogens to challenge a mammal according to known immunization regimens. See, e.g., Hellstrom et al., U.S. Patent 4,918,164. The resulting anti-idiotypic antibodies can be applied, using standard techniques and methods, as: a) vaccines to induce an immune response resulting in protection against the development of Norwalk virus infection; b) immunotherapeutics to induce a patient's immune response to a Norwalk virus infection; c) competing antigens for use in the competitive immunoassay format described above; and d) as purification reagents to separate Norwalk virus-binding substances from non-Norwalk virus-binding substances.

- 19 -

For *in vivo* applications, those skilled in the art will recognize that there are various known techniques to make non-human, e.g., mouse, monoclonal antibodies more compatible with human therapies, e.g., as described in
5 Winter et al., Nature, 349:293-299, 1991; Sahagan et al., J. Immunol., 137:1066-1074, 1986; and Boss et al., U.S. Patent 4,816,397.

It is to be understood that while the invention has been described in conjunction with the detailed
10 description thereof, that the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

- 20 -

CLAIMS:

1. An assay to determine the presence of Norwalk virus or Norwalk virus antigens in a biological sample, the assay comprising the steps of
contacting the sample with a monoclonal antibody
5 that specifically binds to Norwalk virus particles in human stool;
reacting the sample and the monoclonal antibody for a time and under conditions that allow the formation of an immunocomplex between the monoclonal antibody and
10 any Norwalk virus or virus antigens in the sample; and
detecting the immunocomplex, the presence of the immunocomplex indicating the presence of human Norwalk virus or virus antigens in the sample.
2. An assay of claim 1, wherein the immunocomplex
15 is detected by a competitive immunoassay that comprises reacting the monoclonal antibody with the sample and with a competing antigen to which the monoclonal antibody is known to specifically bind.
3. An assay of claim 1, wherein the monoclonal
20 antibody is a first antibody, and the immunocomplex is detected by reaction with a second antibody that is labeled and specifically binds to the first antibody or to human Norwalk virus.
4. An assay of claim 1, wherein the sample is a
25 stool sample.
5. An assay of claim 1, wherein the Norwalk virus antigen is a 58 KDa Norwalk virus protein.

- 21 -

6. An assay to determine the presence of Norwalk virus-specific IgM in a biological sample, the assay comprising the steps of

isolating IgM in the sample;

5 contacting the isolated IgM with purified Norwalk virus or Norwalk virus particles for a time and under conditions that allow the formation of a Norwalk virus-IgM complex between any Norwalk virus-specific IgM in the sample and the purified Norwalk virus or Norwalk virus
10 particles;

contacting the Norwalk virus-IgM complex with a monoclonal antibody that specifically binds to Norwalk virus particles in human stool;

15 reacting the complex and the monoclonal antibody for a time and under conditions that allow the formation of an immunocomplex between the monoclonal antibody and any Norwalk virus-IgM complexes; and

20 detecting the immunocomplex, the presence of the immunocomplex indicating the presence of Norwalk virus-specific IgM in the sample.

7. An assay of claim 6, wherein the IgM is isolated with an anti-human IgM antibody.

8. An assay of claim 6, wherein the sample is a serum sample.

25 9. An assay of claim 1 or 6, wherein the monoclonal antibody specifically binds to a 58 KDa Norwalk virus protein.

10. An assay of claim 1 or 6, wherein the monoclonal antibody is monoclonal antibody 1D8.

- 22 -

11. An assay of claim 1 or 6, wherein the monoclonal antibody is monoclonal antibody 1C9.

12. A monoclonal antibody that specifically binds to a Norwalk virus antigen in stool from a human patient
5 infected with Norwalk virus.

13. A monoclonal antibody of claim 12, wherein the Norwalk virus antigen is a 58 KDa Norwalk virus protein.

14. A monoclonal antibody of claim 12 that is
10 monoclonal antibody 1C9.

15. A monoclonal antibody of claim 12 that is monoclonal antibody 1D8.

16. An anti-idiotypic antibody that specifically binds to the monoclonal antibody of claim 12.

15 17. A vaccine suitable for administration to humans that comprises an anti-idiotypic antibody of claim 16.

18. An immunoassay kit to detect human Norwalk virus in a sample, the kit comprising a monoclonal
20 antibody of claim 12 and means for determining binding of the monoclonal antibody to human Norwalk virus in the sample.

19. An immunoassay kit to detect human Norwalk virus-specific IgM in a sample, the kit comprising a
25 monoclonal antibody of claim 12 and means for determining binding of the monoclonal antibody to human Norwalk virus bound to Norwalk virus-specific IgM in the sample.





FIG. 1



INTERNATIONAL SEARCH REPORT

 International application No.
PCT/US96/13215
A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : Please See Extra Sheet.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, DIALOG, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y	JIANG et al. Expression, Self-Assembly, and Antigenicity of the Norwalk Virus Capsid Protein. Journal of Virology. November 1992, Vol. 66, No. 11, pages 6527-6532, see entire document.	1-16, 18-19 ----- 17
Y	GRAHAM et al. Norwalk Virus Infection of Volunteers: New Insights Based on Improved Assays. The Journal of Infectious Diseases. July 1994, Vol. 170, pages 34-43, see entire document.	1-15, 18, 19
X,P ---- Y,P	HERRMANN et al. Monoclonal Antibodies for Detection of Norwalk Virus Antigen in Stools. Journal of Clinical Microbiology. September 1995, Vol. 33, No. 9, pages 2511-2513, see entire document.	1-16, 18, 19 ----- 17



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

16 SEPTEMBER 1996

Date of mailing of the international search report

24 OCT 1996

 Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JAY WILLIAMS

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/13215

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y	GREEN et al. Comparison of the Reactivities of Baculovirus-Expressed Recombinant Norwalk Virus Capsid Antigen with Those of the Native Norwalk Virus Antigen in Serologic Assays and Some Epidemiologic Observations. Journal of Clinical Microbiology. August 1993, Vol. 31, No. 8, pages 2185-2191, see entire document.	1-16,18,19 ----- 17
A	OKHUYSEN et al. Viral Shedding and Fecal IgA Response after Norwalk Virus Infection. The Journal of Infectious Diseases. March 1995, Vol. 171, pages 566-569, see entire document.	1-19

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/13215

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A61K 39/395, 39/40, 39/42, 38/00; G01N 33/53, 33/537, 33/543, 33/563; C12N 7/00, 7/01; C07K 16/00; C12P 21/08

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

424/147.1, 130.1, 131.1, 134.1, 136.1, 139.1 141.1, 159.1; 530/388.3, 387.2, 388.1, 388.2; 436/513; 435/5, 7.1, 7.92, 7.93, 7.94, 7.95, 235.1

B. FIELDS SEARCHED

Minimum documentation searched

Classification System: U.S.

424/147.1, 130.1, 131.1, 134.1, 136.1, 139.1 141.1, 159.1; 530/388.3, 387.2, 388.1, 388.2; 436/513; 435/5, 7.1, 7.92, 7.93, 7.94, 7.95, 235.1